

**MOHAMED KHIDER UNIVERSITY OF BISKRA.**  
**FACULTY OF EXACT SCIENCES AND NATURAL AND LIFE SCIENCES**  
**DEPARTMENT OF BIOLOGY**

**COURSE TITLE: General Chemistry I**  
**CHAPTER I**  
**Level: 1<sup>st</sup> year LMD**

**Dr: Ismail DAOUD**

**Academic year: 2024/2025**

## 1. Generality


1.1. Atom, nucleus, isotopy,...

1.1. Stability and cohesion of the nucleus, binding energy per nucleon,...

## 1. Generality

Matter is made up of atoms, themselves made up of a nucleus surrounded by an electronic cloud (Fig. 1 and Table. 1). The central spherical nucleus is composed of A nucleons distributed into:

- Z protons (charge  $q = + e$ ; mass  $m_p$ ).
- N neutrons (charge  $q = 0$ ; mass  $m_n$ ).



**Figure:** Schematic view of the structure of matter

### 1.1. Atom, nucleus, isotopy,

#### 1.1.1. Atoms and subatomic particles:

- Atoms are mostly empty space.
- Protons and neutrons occupy small, dense nucleus.
- Electrons move around outside of the nucleus.

The proton and neutron are each more than 1800 times heavier than the electron.

Charge of proton = -(charge of electron) = e

**Table :** Main characteristics of the electron, proton and neutron

Particules and symboles		Autors	Charge*	Masses ( $m_e$ , $m_p$ , $m_n$ )
<b>Electron</b>	<b>e -</b>	J. J. Thomson (1897) R. A. Millikan (1911)	$- 1,6 \cdot 10^{-19} \text{ C}$	$9,11 \cdot 10^{-31} \text{ kg}$
<b>Proton</b>	<b>p</b>	E. Rutherford (1910)	$+ 1,6 \cdot 10^{-19} \text{ C}$	$1,67 \cdot 10^{-27} \text{ kg}$
<b>Neutron</b>	<b>n</b>	J. Chadwick (1932)	0	$1,67 \cdot 10^{-27} \text{ kg}$

### 1.1.2. Nucleus characteristic ${}^A_ZX^q$

The nucleus always has a positive electric charge “q” which is an integer multiple of the elementary charge e:  $q = Z \cdot e$  ( $Z \in \mathbb{N}$ , N integer).

The nucleus is characterized by two numbers Z and A:

Z: is called atomic number: characteristic of the atom, represents the number of electrons.

Z: designates the number of charge: characteristic of the nucleus represents the number of protons.

A: designates the number of a mass, it also represents the number of protons and neutrons constituting the nucleus (the number of nucleons); it is the whole number closest to the atomic mass evaluated in atomic mass unit (u) such that:  $A = (m_p + m_n) - m_{\text{nucleus}}$ .

**Definition:** The atomic mass unit u is equal to 1/12 of the mass of the isotope 12 of carbon  ${}^{12}\text{C}$  (including the mass of the gram atom “atg” of  ${}^{12}\text{C} = 12.00000\text{g}$ ). So :

$$1u = \frac{1}{12} * \frac{12 * 10^{-3}}{N} = \frac{10^{-3}}{6.0221367 * 10^{23}} = 1.6605402 * 10^{-27} \text{Kg}$$

Or:  $N$ : Avogadro number =  $6.022167 * 10^{23} \text{ mol}^{-1}$ .

Avogadro's constant represents the number of particles per mole.

### 1.1.3. Concept of isotopy

Isotopes of the same element are species which have the same number of protons (same Z) and a different number of neutrons (therefore different A).

Example :

Carbon	Hydrogen
${}^{12}_6\text{C}$	${}^1_1\text{H}$
${}^{13}_6\text{C}$	${}^2_1\text{H}$
${}^{14}_6\text{C}$	${}^3_1\text{H}$

#### **a. Relative abundance of different isotopes**

By natural abundance we designate the percentage in number of atoms of each of the isotopes present in the natural mixture.

The natural abundance of each isotope is always the same regardless of the origin of the sample studied.

Example: Carbon has two natural stable isotopes: commonly called Carbon 12 and Carbon 13. Their natural abundances are as follows:

<b>Number of Mass:</b>	<sup>12</sup> C	<sup>13</sup> C
<b>Abundance:</b>	98.9 %	1.1%

**b. Molar Mass of the element (Average or isotopic)**

An element is made up of a mixture of various isotopes and the proportions of these various isotopes are constant. We will be able to define for each element an average molar mass which will take into account its composition.

$$M_A = \frac{\sum_{i=0}^n X_i * M_i}{\sum_{i=0}^n X_i}$$

$X_i$ : designating the natural abundance of isotope  $i$  with molar mass  $M_i$ .

$M_i$ : Molar mass of isotope  $i$ .

Example: Calculate the average isotopic mass of carbon:

$$M_C = 98.89 * M(^{12}\text{C}) + 1.1 * M(^{13}\text{C})/100$$

If we do not need extreme precision we can assimilate the molar masses of each of the isotopes to their mass number.

$$M_C = 98.89 * 12 + 1.1 * 13/100 = 12.02 \text{ g.mol}^{-1}$$

**Note:** For certain elements, there are also unstable natural or artificial isotopes called radioactive. Due to their instability, their abundance varies over time and is therefore never specified.

**1.1.4. Stability and cohesion of the nucleus, binding energy per nucleon**

**a. Mass and energy: Einstein's relationship**

During a nuclear transformation (natural or induced), the mass of the products is always a little lower than the mass of the reactants. The mass loss is denoted  $\Delta m$ . Associated with this loss of mass, there is a release of energy whose value  $\Delta E$  is given by the Einstein relation:

$$\Delta E = (\Delta m) C_0^2$$

$C_0$ : is the speed of light in a vacuum. The convention adopted here is to represent  $\Delta m$  and  $\Delta E$  by positive quantities.

**b. Cohesive energy of a nucleus**

The mass of an atomic nucleus is always a little less than the sum of the masses of its nucleons. The difference  $\Delta m$  between the mass of the nucleons and that of the nucleus is called mass defect. The corresponding quantity of energy  $\Delta E$  given by the relation  $\Delta E = (\Delta m).C_0^2$  is the energy which would be released if the nucleus was formed from nucleons.

A nucleus is all the more stable as this quantity  $\Delta E$  is large. This is why  $\Delta E$  is called cohesion energy (or binding energy) of the nucleus. To compare the stabilities of the nuclei of various elements, we compare the cohesion energies referred to 1 nucleon, and expressed in MeV/nucleon\* (\*:  $1 \text{ eV} = 1.602 \cdot 10^{-19} \text{ J}$ ;  $1 \text{ MeV} = 1.602 \cdot 10^{-19} \times 10^6 = 1.602 \cdot 10^{-13} \text{ J}$ ).